

INFORMATION IN THE TABLE BELOW AND IN THE TABLES ON PAGES 3-5 MAY BE USEFUL IN ANSWERING THE QUESTIONS IN THIS SECTION OF THE EXAMINATION.

DO NOT DETACH FROM BOOK.

PERIODIC TABLE OF THE ELEMENTS

H	1.008	4
Li	6.94	Be
11	12	
Na	22.99	Mg
19	20	21
K	39.10	Ca
37	38	39
Rb	85.47	Sr
55	56	57
Cs	132.91	Ba
87	88	89
Fr	(223)	[†] Ac
137.33	138.91	178.49
226.02	227.03	(261)

H	2	He
		4.00
B	5	6
10.81	12.01	C
13	14	N
Al	Si	O
26.98	28.09	F
30	31	Ne
Zn	Ga	16.00
65.39	69.72	19.00
46	47	20.18
Pd	Ag	17
102.91	106.42	18
48	49	17
Cd	In	17
107.87	112.41	17
50	51	17
Sn	Sb	17
114.82	118.71	17
53	52	17
Te	I	17
121.75	127.60	17
80	81	17
Tl	Pb	17
195.08	196.97	17
204.38	207.2	17
208.98	(209)	17
(210)	(221)	17

Ce	58	59	60	61	62	63	64	65	66	67	68	69	70	71
140.12	140.91	144.24	(145)	150.4	151.97	157.25	158.93	162.50	164.93	167.26	168.93	173.04	174.97	
90	91	92	93	94	95	96	97	98	99	100	101	102	103	
Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr	
232.04	231.04	238.03	(237)	(244)	(243)	(247)	(247)	(251)	(252)	(258)	(259)	(262)	(262)	

*Lanthanide Series

†Actinide Series

GO ON TO THE NEXT PAGE.

STANDARD REDUCTION POTENTIALS IN AQUEOUS SOLUTION AT 25°C

	Half-reaction	$E^\circ(V)$
$\text{F}_2(g) + 2e^-$	$\rightarrow 2\text{F}^-$	2.87
$\text{Co}^{3+} + e^-$	$\rightarrow \text{Co}^{2+}$	1.82
$\text{Au}^{3+} + 3e^-$	$\rightarrow \text{Au}(s)$	1.50
$\text{Cl}_2(g) + 2e^-$	$\rightarrow 2\text{Cl}^-$	1.36
$\text{O}_2(g) + 4\text{H}^+ + 4e^-$	$\rightarrow 2\text{H}_2\text{O}(l)$	1.23
$\text{Br}_2(l) + 2e^-$	$\rightarrow 2\text{Br}^-$	1.07
$2\text{Hg}^{2+} + 2e^-$	$\rightarrow \text{Hg}_2^{2+}$	0.92
$\text{Hg}^{2+} + 2e^-$	$\rightarrow \text{Hg}(l)$	0.85
$\text{Ag}^+ + e^-$	$\rightarrow \text{Ag}(s)$	0.80
$\text{Hg}_2^{2+} + 2e^-$	$\rightarrow 2\text{Hg}(l)$	0.79
$\text{Fe}^{3+} + e^-$	$\rightarrow \text{Fe}^{2+}$	0.77
$\text{I}_2(s) + 2e^-$	$\rightarrow 2\text{I}^-$	0.53
$\text{Cu}^+ + e^-$	$\rightarrow \text{Cu}(s)$	0.52
$\text{Cu}^{2+} + 2e^-$	$\rightarrow \text{Cu}(s)$	0.34
$\text{Cu}^{2+} + e^-$	$\rightarrow \text{Cu}^+$	0.15
$\text{Sn}^{4+} + 2e^-$	$\rightarrow \text{Sn}^{2+}$	0.15
$\text{S}(s) + 2\text{H}^+ + 2e^-$	$\rightarrow \text{H}_2\text{S}(g)$	0.14
$2\text{H}^+ + 2e^-$	$\rightarrow \text{H}_2(g)$	0.00
$\text{Pb}^{2+} + 2e^-$	$\rightarrow \text{Pb}(s)$	-0.13
$\text{Sn}^{2+} + 2e^-$	$\rightarrow \text{Sn}(s)$	-0.14
$\text{Ni}^{2+} + 2e^-$	$\rightarrow \text{Ni}(s)$	-0.25
$\text{Co}^{2+} + 2e^-$	$\rightarrow \text{Co}(s)$	-0.28
$\text{Cd}^{2+} + 2e^-$	$\rightarrow \text{Cd}(s)$	-0.40
$\text{Cr}^{3+} + e^-$	$\rightarrow \text{Cr}^{2+}$	-0.41
$\text{Fe}^{2+} + 2e^-$	$\rightarrow \text{Fe}(s)$	-0.44
$\text{Cr}^{3+} + 3e^-$	$\rightarrow \text{Cr}(s)$	-0.74
$\text{Zn}^{2+} + 2e^-$	$\rightarrow \text{Zn}(s)$	-0.76
$2\text{H}_2\text{O}(l) + 2e^-$	$\rightarrow \text{H}_2(g) + 2\text{OH}^-$	-0.83
$\text{Mn}^{2+} + 2e^-$	$\rightarrow \text{Mn}(s)$	-1.18
$\text{Al}^{3+} + 3e^-$	$\rightarrow \text{Al}(s)$	-1.66
$\text{Be}^{2+} + 2e^-$	$\rightarrow \text{Be}(s)$	-1.70
$\text{Mg}^{2+} + 2e^-$	$\rightarrow \text{Mg}(s)$	-2.37
$\text{Na}^+ + e^-$	$\rightarrow \text{Na}(s)$	-2.71
$\text{Ca}^{2+} + 2e^-$	$\rightarrow \text{Ca}(s)$	-2.87
$\text{Sr}^{2+} + 2e^-$	$\rightarrow \text{Sr}(s)$	-2.89
$\text{Ba}^{2+} + 2e^-$	$\rightarrow \text{Ba}(s)$	-2.90
$\text{Rb}^+ + e^-$	$\rightarrow \text{Rb}(s)$	-2.92
$\text{K}^+ + e^-$	$\rightarrow \text{K}(s)$	-2.92
$\text{Cs}^+ + e^-$	$\rightarrow \text{Cs}(s)$	-2.92
$\text{Li}^+ + e^-$	$\rightarrow \text{Li}(s)$	-3.05

GO ON TO THE NEXT PAGE.

ADVANCED PLACEMENT CHEMISTRY EQUATIONS AND CONSTANTS

ATOMIC STRUCTURE

$$\begin{aligned}E &= h\nu & c &= \lambda\nu \\ \lambda &= \frac{h}{mv} & p &= mv \\ E_n &= \frac{-2.178 \times 10^{-18}}{n^2} \text{ joule}\end{aligned}$$

EQUILIBRIUM

$$K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$$

$$K_b = \frac{[\text{OH}^-][\text{HB}^+]}{[\text{B}]}$$

$$\begin{aligned}K_w &= [\text{OH}^-][\text{H}^+] = 1.0 \times 10^{-14} @ 25^\circ\text{C} \\ &= K_a \times K_b\end{aligned}$$

$$\begin{aligned}\text{pH} &= -\log[\text{H}^+], \text{pOH} = -\log[\text{OH}^-] \\ 14 &= \text{pH} + \text{pOH}\end{aligned}$$

$$\text{pH} = \text{p}K_a + \log \frac{[\text{A}^-]}{[\text{HA}]}$$

$$\text{pOH} = \text{p}K_b + \log \frac{[\text{HB}^+]}{[\text{B}]}$$

$$\text{p}K_a = -\log K_a, \text{p}K_b = -\log K_b$$

$$K_p = K_c (RT)^{\Delta n},$$

where Δn = moles product gas – moles reactant gas

THERMOCHEMISTRY/KINETICS

$$\Delta S^\circ = \sum S^\circ \text{ products} - \sum S^\circ \text{ reactants}$$

$$\Delta H^\circ = \sum \Delta H_f^\circ \text{ products} - \sum \Delta H_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \sum \Delta G_f^\circ \text{ products} - \sum \Delta G_f^\circ \text{ reactants}$$

$$\begin{aligned}\Delta G^\circ &= \Delta H^\circ - T\Delta S^\circ \\ &= -RT \ln K = -2.303 RT \log K \\ &= -n\mathcal{F}E^\circ\end{aligned}$$

$$\Delta G = \Delta G^\circ + RT \ln Q = \Delta G^\circ + 2.303 RT \log Q$$

$$q = mc\Delta T$$

$$C_p = \frac{\Delta H}{\Delta T}$$

$$\ln[\text{A}]_t - \ln[\text{A}]_0 = -kt$$

$$\frac{1}{[\text{A}]_t} - \frac{1}{[\text{A}]_0} = kt$$

$$\ln k = \frac{-E_a}{R} \left(\frac{1}{T} \right) + \ln A$$

E = energy	v = velocity
ν = frequency	n = principal quantum number
λ = wavelength	m = mass
p = momentum	

$$\text{Speed of light, } c = 3.0 \times 10^8 \text{ m s}^{-1}$$

$$\text{Planck's constant, } h = 6.63 \times 10^{-34} \text{ J s}$$

$$\text{Boltzmann's constant, } k = 1.38 \times 10^{-23} \text{ J K}^{-1}$$

$$\text{Avogadro's number} = 6.022 \times 10^{23} \text{ mol}^{-1}$$

$$\text{Electron charge, } e = -1.602 \times 10^{-19} \text{ coulomb}$$

$$1 \text{ electron volt per atom} = 96.5 \text{ kJ mol}^{-1}$$

Equilibrium Constants

K_a (weak acid)

K_b (weak base)

K_w (water)

K_p (gas pressure)

K_c (molar concentrations)

S° = standard entropy

H° = standard enthalpy

G° = standard free energy

E° = standard reduction potential

T = temperature

n = moles

m = mass

q = heat

c = specific heat capacity

C_p = molar heat capacity at constant pressure

E_a = activation energy

k = rate constant

A = frequency factor

Faraday's constant, $\mathcal{F} = 96,500$ coulombs per mole of electrons

Gas constant, $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$

$= 0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1}$

$= 62.4 \text{ L torr mol}^{-1} \text{ K}^{-1}$

$= 8.31 \text{ volt coulomb mol}^{-1} \text{ K}^{-1}$

GO ON TO THE NEXT PAGE.

GASES, LIQUIDS, AND SOLUTIONS

$$PV = nRT$$

$$\left(P + \frac{n^2 a}{V^2} \right) (V - nb) = nRT$$

$$P_A = P_{total} \times X_A, \text{ where } X_A = \frac{\text{moles A}}{\text{total moles}}$$

$$P_{total} = P_A + P_B + P_C + \dots$$

$$n = \frac{m}{M}$$

$$K = {}^\circ\text{C} + 273$$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$D = \frac{m}{V}$$

$$u_{rms} = \sqrt{\frac{3kT}{m}} = \sqrt{\frac{3RT}{M}}$$

$$KE \text{ per molecule} = \frac{1}{2}mv^2$$

$$KE \text{ per mole} = \frac{3}{2}RT$$

$$\frac{r_1}{r_2} = \sqrt{\frac{M_2}{M_1}}$$

molarity, M = moles solute per liter solution

molality = moles solute per kilogram solvent

$$\Delta T_f = iK_f \times \text{molality}$$

$$\Delta T_b = iK_b \times \text{molality}$$

$$\pi = iMRT$$

$$A = abc$$

P = pressure

V = volume

T = temperature

n = number of moles

D = density

m = mass

v = velocity

u_{rms} = root-mean-square speed

KE = kinetic energy

r = rate of effusion

M = molar mass

π = osmotic pressure

i = van't Hoff factor

K_f = molal freezing-point depression constant

K_b = molal boiling-point elevation constant

A = absorbance

a = molar absorptivity

b = path length

c = concentration

Q = reaction quotient

I = current (amperes)

q = charge (coulombs)

t = time (seconds)

E° = standard reduction potential

K = equilibrium constant

OXIDATION-REDUCTION; ELECTROCHEMISTRY

$$Q = \frac{[C]^c [D]^d}{[A]^a [B]^b}, \text{ where } aA + bB \rightarrow cC + dD$$

$$I = \frac{q}{t}$$

$$E_{cell} = E_{cell}^\circ - \frac{RT}{nF} \ln Q = E_{cell}^\circ - \frac{0.0592}{n} \log Q @ 25^\circ\text{C}$$

$$\log K = \frac{nE^\circ}{0.0592}$$

Gas constant, $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$

= $0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1}$

= $62.4 \text{ L torr mol}^{-1} \text{ K}^{-1}$

= $8.31 \text{ volt coulomb mol}^{-1} \text{ K}^{-1}$

Boltzmann's constant, $k = 1.38 \times 10^{-23} \text{ J K}^{-1}$

K_f for $\text{H}_2\text{O} = 1.86 \text{ K kg mol}^{-1}$

K_b for $\text{H}_2\text{O} = 0.512 \text{ K kg mol}^{-1}$

1 atm = 760 mm Hg

= 760 torr

STP = 0.00°C and 1.0 atm

Faraday's constant, $F = 96,500 \text{ coulombs per mole of electrons}$

GO ON TO THE NEXT PAGE.